

PHTHALIC ACID ESTERS AND POLYCYCLIC AROMATIC HYDROCARBONS IN HOUSEHOLD DUST

E. D. Vicente¹, A. Vicente¹, T. Nunes¹, A. Calvo², C. del Blanco-Alegre², F. Oduber², A. Castro², R. Fraile², F. Amato³, X. Querol³, C. Alves¹

1 INTRODUCTION

Household dust is a complex mixture of particles of both indoor and outdoor origin, including organic, inorganic and biological particles (Naspinski et al., 2008). Its composition depends on numerous conditions, such as environmental and seasonal factors, ventilation and air filtration, homeowner activities, and in- and outdoor sources (Maertens et al., 2004). Residential dust is recognized as a major source of environmental contaminants, including polycyclic aromatic hydrocarbons (PAHs) and phthalic acid esters (PAEs) (Roberts et al., 2009). The selection of appropriate techniques to assess household dust amount and composition is a major challenge since several different methodologies have been employed. The methodologies used for indoor dust collection include passive (dust settling) and active techniques (surface wiping, press sampling, sweeping, or vacuuming) (Lioy et al., 2002; Morawska and Salthammer, 2003). The aim of the present study was to assess the variation of the PM₁₀ fraction of household dust in chemical composition within a home and in homes with different outdoor surrounding. An active sampling methodology was applied to collect the deposited PM₁₀ fraction directly from the floor.


2 METHODOLOGY

Sampling Sites

4 Houses


House 1	Suburban two-story house with well ventilated kitchen. Two occupants.
House 2	Single story apartment located in the city center. Two occupants.
House 3	Rural two-story house with open fireplace in the living room. Two occupants.
House 4	Single story apartment with small kitchen open to the living room. One occupant.

3 Rooms



Two Sampling Periods

Sampling Methodology



Analysis




Thermo-optical method:
Total Carbon (TC)



GC-MS: PAHs and PAEs

3 RESULTS

1st Sampling



2nd Sampling




Fig. 1. PAEs mass fractions (ng µg⁻¹) in PM₁₀ dust samples collected in two different sampling periods





Fig. 2. PAHs mass fractions (ng µg⁻¹) in PM₁₀ dust samples collected in two different sampling periods.





Fig. 3. PAEs mass fractions (ng µg⁻¹) in PM₁₀ dust samples collected in two different sampling periods.

4 CONCLUSIONS

- Total carbon contribution to the PM₁₀ mass showed temporal and spatial variability ranging from 9.3 to 75%wt.
- Bis(2-ethylhexyl) phthalate (DEHP, n.d.–9.42 ng µg⁻¹ PM₁₀) and di-n-butyl phthalate (DnBP, 0.00–10.2 ng µg⁻¹ PM₁₀) were the major phthalates in the household dust. DnBP has been reported to be largely present in cosmetic and personal care products (Koniecki et al., 2011), while DEHP was the most abundant phthalate compound found in food products and packaging materials (Fierens et al., 2012).
- PAHs had a smaller fractional contribution to indoor dust (Σ₂₀PAH 0.005–0.411 ng µg⁻¹ PM₁₀). The main PAHs contributing to the household dust mass were pyrene (n.d.–0.089 ng µg⁻¹ PM₁₀) and retene (n.d.–0.082 ng µg⁻¹ PM₁₀).
- Although the highest dust loads were recorded for rugs, PAEs and PAHs had the lowest contributions to the total dust levels compared to the ones recorded in hard floorings.
- Although this study is based on small number of samples, the findings underline the importance of organic pollutants accumulated in indoor dust. Larger studies, covering more homes, are needed in order to better understand the indoor dynamics of these pollutants.

ACKNOWLEDGEMENTS

Thanks are given to the Portuguese Foundation of Science and Technology (FCT) and POHP/FSE funding program for the fellowship SFRH/BD/117993/2016. Ana Vicente is funded by national funds (OE), through FCT, I.P., in the scope of the framework contract foreseen in the numbers 4, 5 and 6 of the article 23, of the Decree-Law 57/2016, of August 29, changed by Law 57/2017, of July 19. This work was also financially supported by the project POCI-01-0145-FEDER-029574 funded by FEDER, through COMPETE2020 - Programa Operacional Competitividade e Internacionalização (POCI), and by OE, through FCT/MCTES.

REFERENCES

- Fierens, T., Servaes, K., Van Holderbeke, M., Geerts, L., De Henauw, S., Sioen, I., Vanermen, G. (2012). Food Chem. Toxicol., 50, 2575–2583.
- Koniecki, D., Wang, R., Moody, R.P., Zhu, J. (2011). Environ. Res., 111, 329–336.
- Maertens, R.M., Bailey, J., White, P.A. (2004). Mutat. Res. - Rev. Mutat. Res., 567, 401–425.
- Naspinski, C., Lingenfelter, R., He, L.Y., Cizmas, L., Naufal, Z., Islamzadeh, A., Li, Z., Li, Z., Donnelly, K.C., McDonald, T. (2008). Environ. Int., 34, 988–993.
- Roberts, J.W., Wallace, L.A., Camann, D.E., Dickey, P., Gilbert, S.G., Lewis, R.G., Takaro T.K., in: Rev. Environ. Contam. Toxicol., (2009) pp. 1–39.