

## Winter air quality in an urban area with high coal consumption in domestic devices

E. D. Vicente<sup>1,2</sup>, C. Gonçalves<sup>1</sup>, A. I. Calvo<sup>1</sup>, C. Blanco-Alegre<sup>1</sup>, A. Sánchez de la Campa<sup>3,4</sup>, T. Sainnokhoi<sup>4</sup>, N. Kováts<sup>4</sup>, F. Giardi<sup>5</sup>, S. Nava<sup>5</sup>, G. Calzolai<sup>5</sup> and R. Fraile<sup>1</sup>

<sup>1</sup>Department of Physics, University of León, León, 24071, Spain

<sup>2</sup>Department of Environment and Planning, Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, 3810-193, Portugal

<sup>3</sup>Associate Unit CSIC-University of Huelva "Atmospheric Pollution", Centre for Research in Sustainable Chemistry - CIQSO, University of Huelva, Huelva, E21071, Spain

<sup>4</sup>Centre for Environmental Sciences, University of Pannonia, Veszprém, 8200, Hungary

<sup>5</sup>INFN - Firenze, National Institute for Nuclear Physics - Florence division, 50019, Sesto Fiorentino, Italy

Keywords: air quality, PM<sub>10</sub>, PM-bound constituents, toxicity.

Associated conference topics: 2.9, 2.11

Presenting author email: rfral@unileon.es

Air pollution is the 4<sup>th</sup> leading risk factor for mortality worldwide (HEI, 2020). Particulate matter (PM), classified as carcinogenic for humans (Group 1) (Loomis et al. 2013), is the leading environmental contributor to the global burden of disease contributing (European Environment Agency, 2019). Despite the improvements on outdoor air quality in Europe, PM levels can be heightened under episodic events provoked, for example, by seasonal emission sources under stagnant weather conditions. In this study, a winter monitoring campaign was carried out at the city center of León (Spain) in order to evaluate the PM<sub>10</sub> concentrations, chemical composition and toxicity.

A sampling campaign was carried out from 17<sup>th</sup> to 24<sup>th</sup> January to evaluate the PM levels in the city centre of León, during winter, when high usage of domestic heating devices is expected. The monitoring week was one of the coldest of the 2021 – 2022 winter. León is surrounded by mountain ranges on the west, north, and east. The climate is characterized by cold winters and hot summers. León is located in a coal-mining region, where the use of coal in domestic heating devices is still widespread. The main particle emission sources in the urban area is road traffic as well as coal and biomass burning in winter (Blanco-Alegre et al. 2019). Samplers were located on the rooftop of an official building located in front of a circular square with heavy traffic. Meteorological data (wind, temperature, precipitation and relative humidity) was recorded with a weather station. PM<sub>10</sub> samples were collected using a low-volume sampler (TCR Tecora, model 2.004.01) and a high-volume sampler (MCV, model CAV-A/mb) equipped with teflon and quartz fibre filters, respectively. Samples were taken in parallel for 23h30 every day for one week, starting at 1100 and ending at 1030 UTC. Chemical composition was obtained by PIXE (elements), ion chromatography (water-soluble inorganic ions) and thermal-optical analysis (organic and elemental carbon). The ecotoxicity of the PM<sub>10</sub> samples was evaluated by the *Vibrio fischeri* bioluminescence inhibition assay (Vicente et al., 2021).

The average daily temperature recorded during the sampling period was  $3.5 \pm 1.7$  °C and the relative

humidity  $63.7 \pm 4.57\%$ . No precipitation events occurred during the studied period. The daily average PM<sub>10</sub> concentrations were always below the EU daily limit value, ranging between 20.0 and 39.2  $\mu\text{g m}^{-3}$ . OC and EC represented  $22.2\% \pm 2.21\%$  wt. and  $2.69\% \pm 1.69\%$  wt. PM<sub>10</sub>, respectively. Water-soluble ions accounted for 11.1% - 36.7% wt. of the PM<sub>10</sub> mass. The dominant water-soluble species were sulphate, nitrate and ammonium which globally represented 64% - 76% of the total ionic content. These secondary inorganic ions were significantly correlated. The dominant elements in PM<sub>10</sub> samples were Si (2.66% – 6.71% wt. PM<sub>10</sub>) and Cl (3.59 – 5.18% wt. PM<sub>10</sub>). The concentrations of Se, a coal burning tracer, ranged from 2.03 to 10.6  $\text{ng m}^{-3}$ . All the samples inhibited the bioluminescence of the bacteria, indicating that all the samples were toxic.

This work was partially supported by the Junta de Castilla y Leon co-financed with European FEDER funds (Grant LE025P20) and by the AEROHEALTH project (Ministry of Science and Innovation, co-financed with European FEDER funds, grant PID2019-106164RBI00). This research has also been supported by the RADIATE project under the Grant Agreement 824096 from the EU Research and Innovation programme HORIZON 2020. Furthermore, it is part of the project TED2021-132292B-I00, funded by MCIN/AEI/10.13039/501100011033 and by the European Union "NextGenerationEU"/PRTR. The financial support to CESAM by FCT/MCTES (UIDP/50017/2020+UIDB/50017/2020).

Blanco-Alegre C, Calvo AI, Coz E, et al (2019) Environ. Pollut. **246**, 336–345.

European Environment Agency (2019) Health impacts of air pollution

HEI (2020) State of Global Air 2020. Special Report. Health Effects Institute, Boston, MA

Loomis D, Grosse Y, Lauby-Secretan B, et al (2013) Lancet Oncol. **14**, 1262–1263.

Vicente, E.D., Figueiredo, D., Gonçalves, C., Lopes, I., Oliveira, H., Kováts, N., Pinheiro, T., Alves, C.A. (2021) Sci. Total Environ. **782**, 146820.